References:

1] International Conferences on Harmonization (2008), Draft Revised Guidance on Impurities in New Drug Substances. Q3A (R2).

2] International Conferences on Harmonization (2003), Draft Revised Guidance on Impurities in New Drug Products. Q3B (R2).

3] International Conferences on Harmonization, Impurities Guidelines for residual solvents. Q3C (R5)

4] AhujaS. (1998),*Impurities Evaluation of Pharmaceuticals*. NY, Marcel Dekker.

5] GorogS.(2000),*Identification and Determination of Impurities in Drugs*. Amsterdam.

6] RajN.D., AnbazhaganS., Kunapareddy A., Sunkara N., Chusena N., (2012), Validated stability indicating gradient RP-HPLC method for the estimation of antihypertensive drugs in bulk and pharmaceutical dosage.,*International Current Pharmaceutical Journal,***1(11),**336-341

7] Hamrapurkar P.D. and Gadapayale K. K.,(2013), Optimization and Validation of RP - HPLC Stability Indicating Method for determination of Olmesartan Medoxomil and Its degraded product.,*Int. Journal of Applied Science and Engineering.,* **11(2),**137-147

8] Saminathan J. and Vetrichelvan T., (2011), Method development and validation of Olmesartan, Amlodipine and Hydrochlorothiazide in combined tablet dosage, *Int. Journal of Pharma. Research & Analysis*, **1(1),** 7-14.

9] Zaveri M., Dhru B., Parmar J., Brahmbhatt.V, khandhar A,(2012)Simultaneous estimation of Olmesartan medoxomil and Hydrochlorothiazide by validated reversed phase high performance liquid chromatography., *Int.journal of pharmaceutical research and bio-science,***1 (1)**

10] RajaB. and Lakshmana RaoA.,(2011),Development and validation of a reversed phase hplc method for simultaneous estimation of Olmesartan and Hydrochlorothiazide in combined tablet dosage form.,*Inte. journal of research in pharmacy and chemistry,***1(3)**

11] GodseV.P., BhosaleA.V., BafanaY.S,Borkar D.D, (2010), ICH guidance in practice: Validated stability-indicating HPLC method for simultaneous determination of Olmesartan medoximil and Hydrochlorothiazide in combination drug products. *J. Anal. Chem.***5(2),**137-144

12] SharmaR.N., PancholiS.S., (2010) , RP-HPLC-DAD method for determination of Olmesartan medoxomil in bulk and tablets exposed to forced conditions., *Acta Pharm.***60(1),**13-24

13] DoshiN., ShethA., Patel C.N., (2012), Validated revered phase high performance liquid chromatographic method for simultaneous estimation of Olmesartan medoximil and Hydrochlorothiazide and Amlodipine besylate in newly designed pharmaceutical dosage form*International research journal of pharmacy*, **3(11)**

14] Ashok kumarJ., SathyaA., Senthil kumarK., Patil S. N., PrathapB., Lokes S.B., GopalV., (2010), Simultaneous Estimation of Olmesartan medoxomil and Hydrochlorothiazide by RP-HPLC Method from Combined Dosage Forms ., *Int. J. Res. Pharm. Sci.***1(1),** 24-27

15] ParikhP., SahooU., ZanvarA, SethA.K.,(2013),Derivative spectrophotometric method for simultaneous estimation of Chlorthalidone and olmesartan medoxomil in their tablet dosage form., *International journal of pharmaceutical sciences,***4(4),**1

16] PanchumarthyR., GarikapatiD., (2013),Improved Rapid HPLC Method for the Separation of Five Anti-Hypertensive Agents Using C18 Column: Application to Hydrochlorothiazide Determination in Bulk and Tablet Dosage., *IOSR Journal Of Pharmacy,***3(5),** 07-19

17] SivasakthiR.., Anudeepa J., Senthil kumarC., RamyaR.., RajendranS.S. , MoorthiC., Venkatnarayanan, (2011),Development and validation of RP-HPLC and UV-Spectrophotometric method for Olmesartan medoxomil and Hydrochlorothiazide in combined dosageform.,*Scholars Research Library*, **3(3),** 478-484**.**

18] PandyaG.P. and JoshiH.S., (2013),Development and validation of stability indicating HPLC assay method forsimultaneous determination of Amlodipine

besylate, Olmesartan medoxomiland Hydrochlorothiazide in tablet formulation.,*Pelagia Research Library,***4(2),**145-152**.**

19] Ashutosh KumarS., Debnath M., J. V. L. N. Seshagiri Rao and GowriSankarD., (2014), A new and rapid analytical method development & validation for Simultaneous estimation of Hydrochlorothiazide, Amlodipine &Olmesartan in tablet dosage form by using RP-HPLC.,*Journal of Chemical and Pharmaceutical Research,***6(5),** 1208-1213.

20] BojjaS., ThimmarajuM. and Nerella R.,(2012), Simultaneous determination of Tenofovir disoproxil fumarate and Lamivudine by UV Spectrophotometric Method.,*International Journal of Pharmacy and Pharma Science Research*, **2(1)** 9-15

21] Murugan S., Sikdar P., Debnath S., Niranjan Babu M, (2013),Development and validation of first order derivative uv spectrophotometric method for the estimation of Tenofovir disoproxil fumarate, Lamivudine and Efavirenz in bulk and tablet dosage form.,*Inte. Journal of Pharma Research &Analysis,***3(1) ,**29-32.

22] Appala RajuN. and BegumS., (2008),Simultaneous RP-HPLC Method for the Estimation of the Emtricitabine, Tenofovir Disoproxil Fumerate and Efavirenz in Tablet Dosage Forms.,*Research J. Pharm. and Tech.***1(4)**.

23] DevanaboyinaN., Satyanarayana T. and Ganga Rao B.,(2012), Hplc method development and validation for simultaneous estimation of Tenofovir and Emtricitabine in combined pharmaceutical dosage form*InteJournal of Research in Pharma and Biomedical Sciences,* **3 (1)**

24] DhaneshwarS. R., Havele S.,(2012),Development and validation of a stability-indicating LC method for the determination of Tenofovir Disoproxil Fumarate in pharmaceutical formulation*J. Sci. Technol*.**34 (6),** 615-622.

25] Kavita K.Y, Getta.G,Hariprasad R., VenkatnarayanaR., Kaviarasu M., (2013), Development and validation of RP-UPLC analytical method for simultaneous estimation of Emtricitabine, Rilpivirine, Tenofovir disoproxil fumarate and its pharmaceutical dosage forms., *International research journal of pharmacy*,**4(1)**.

26] HussenS.S., ShenoyP., Udupa And Muddu Krishna, (2013),Development and validation of stability indicating rp-hplc method for Tenofovir nanoparticle formulation.,*International Journal of Pharmacy and Pharmaceutical Sciences*, **5(2)**.

27] AbdelHay M. H., GazyA.A, ShaalanR.A., and AshourH.K., (2013), Simple Spectrophotometric Methods for Determination of Tenofovir Fumarate and Emtricitabine in Bulk Powder and in Tablets.,*Hindawi Publishing Corporation Journal of Spectroscopy* .

28] Komaroju D., Nagarjuna ReddyG., DhanalakshmiK., (2013),Method Development and Validation for Simultaneous Estimation of Emtricitabine and Tenofovir Disoproxil Fumarate in Pure and Tablet Dosage Form by using RP-HPLC was developed.,*International Journal of Pharma Research & Review*,**2(10),**1-11

29] Kumar Pradeep, Dwivedi S.C. and KushnoorA., (2011), Validated HPTLC Method for the Determination of Tenofoviras Bulk Drug and in Pharmaceutical Dosage Form.,*Research Journal of Chemical Sciences,***1(7),** 33-37.

30] Santhosam D., Bhavani N.V., SureshM., (2012), Development and validation of rp-hplc method for the simultaneous estimation of Emtricitabine and Tenofovir from bulk and dosage form.,*International journal of pharmaceutical chemistry research,***1(4),**2278

31] Venkateswara raoJ., Appala rajuN., , vanitha K.P., Mukkanti K. and srinivasu K., (2008), Simultaneous estimation of Tenofovir Disoproxil, Emtricitabine and Efavirenz in tablet dosage form by RP-HPLC.,*orient. j. chem.*, **24(2),** 645-650

32] Soni A. and Thakral S., (2013), Simultaneous Estimation of Tenofovir and Emtricitabine in Human Plasma Using HPLC after Protein Precipitation Extraction.,*J Anal Bioanal Tech*, **4(3)**

33] NevaseP.A.,NimjeH.M., OswalR.J., AntreR.V., KshirsagarS.S., (2011), UV spectrophotometric method for estimation of Tenofovir Disoproxil Fumarate tablet dosage form ., *IJPRD*, **3(3),**73 - 75

34] Validation of Analytical Procedures: Text and Methodology Q2 (R1). Federal Register, March 1995.

35] Validation of Analytical Procedures: Methodology Q2 (R1). Federal Register, May 1997.